

STATISTICS OF VELOCITY FROM SPECTRAL DATA: MODIFIED VELOCITY CENTROIDS

A. LAZARIAN¹ AND A. ESQUIVEL¹

Draft version February 2, 2008

ABSTRACT

We address the problem of studying interstellar turbulence using spectral line data. We find a criterion when the velocity centroids may provide trustworthy velocity statistics. To enhance the scope of centroids applications, we construct a measure that we term “modified velocity centroids” (MVCs) and derive an analytical solution that relates the 2D spectra of the modified centroids with the underlying 3D velocity spectrum. We test our results using synthetic maps constructed with data obtained through simulations of compressible magnetohydrodynamical (MHD) turbulence. We show that the modified velocity centroids (MVCs) are complementary to the Velocity Channel Analysis (VCA) technique. Employed together, they make determining of the velocity spectral index more reliable and for wider variety of astrophysical situations.

Subject headings: turbulence – ISM: general, structure – MHD – radio lines: ISM.

1. INTRODUCTION

The interstellar medium (ISM) is turbulent, as known from observations of non-thermal cloud velocities (Larson 1992), and also expected on theoretical grounds, because of the very large Reynolds numbers (defined as the ratio of inertial to viscous forces). Understanding turbulence is crucial for understanding of many physical processes, including energy injection into the ISM, non-photoelectric heating of the ISM, star formation, propagation of cosmic rays, and heat transport. (see review by Vázquez-Semadeni et al. 2000, Cho & Lazarian 2003a), and references therein.

Obtaining properties of ISM turbulence from observations is a long-standing problem (see review by Lazarian 1999, and references therein). The statistics of the random velocity field is essential for describing the turbulence. To obtain the velocity statistics, so-called velocity centroids (Munch 1958) have been frequently attempted. However, the separation of the velocity and density contributions to the velocity centroids has always been a problem. Therefore the relation between the statistics of velocity and velocity centroids is frequently claimed to be trustworthy only when the density fluctuations are negligible (see Vestuto, Ostriker & Stone 2003).

To remedy this situation a statistical technique termed Velocity Channel Analysis, or in short VCA, was developed in Lazarian & Pogosyan (2000). It has been successfully used to obtain the spectra of turbulence in H I in Small Magellanic Cloud and the Milky Way (see Stanimirović & Lazarian 2001, Deshpande, Dwarakanath, & Goss 2000). However, the VCA requires turbulence to be supersonic and to follow power laws (see Esquivel et al. 2003).

This paper will identify a technique that can be reliably used with turbulence even if the turbulence is subsonic and/or does not obey a power-law. The latter case can occur when, for instance, self-gravity modifies the turbulence on small scales. In pursuing this goal we introduce modified velocity centroids (MVCs) and derive an ana-

lytical relation between the statistics of MVCs and the underlying statistics of the 3D turbulent velocity.

2. MODEL: TURBULENCE IN MAGNETIZED GAS

There has been substantial progress in understanding of compressible magnetohydrodynamical (MHD) turbulence (see a review by Cho & Lazarian 2003a) that will guide us in adopting the model for this paper.

Numerical simulations in Cho & Lazarian (2002, 2003b) revealed anisotropic spectra of Alfvén and slow waves, as well as an isotropic spectrum of the fast waves. The anisotropies are elongated along the local direction of magnetic field. However, since this local direction changes from one place to another, the anisotropy in the system of reference related to the mean magnetic field is rather modest, i.e. of the order of $\delta B/B$, where δB is the dispersion of the random magnetic field. For typical ISM $\delta B \sim B$ (Spoelstra 1972) and the expected anisotropy is modest. Therefore it is possible to characterize the turbulence using isotropic statistics (see testing in Esquivel et al. 2003 and in §4).

Only z-components of the velocity are available through Doppler shift measurements. In what follows we shall assume that the velocity and density of the gas can be presented as sums of a mean value and a fluctuating part: $V_z = v_0 + v$, $\rho_{tot} = \rho_0 + \rho$, where $v_0 \equiv \langle V_z \rangle$, $\rho_0 \equiv \langle \rho_{tot} \rangle$. The fluctuating components v and ρ satisfy the condition $\langle v \rangle = 0$, $\langle \rho \rangle = 0$. Henceforth we use $\langle \dots \rangle$ to denote ensemble averaging. The correlation function of the z-component of velocities at two points $\mathbf{x}_1$ and $\mathbf{x}_2$, separated a distance r (with $\mathbf{r} = \mathbf{x}_1 - \mathbf{x}_2$), denoted by subscripts 1 and 2 ($V_{z,1} = v_0 + v_1$, $V_{z,2} = v_0 + v_2$) is $\langle v_1 v_2 \rangle \equiv B_{zz}(\mathbf{r})$. And it is related to the component of the 3D spectrum through a Fourier transform (see Monin & Yaglom 1975):

$$B_{zz}(\mathbf{r}) = \int e^{i\mathbf{k}\cdot\mathbf{r}} F_{zz}(\mathbf{k}) d\mathbf{k}, \quad (1)$$

where

$$F_{zz}(\mathbf{k}) = (F_{LL}(k) - F_{NN}(k)) k_z^2/k^2 + F_{NN}(k) \quad (2)$$

is the projection of the spectral tensor, expressed through the transversal F_{NN} and longitudinal F_{LL} (normal and

¹ Astronomy Department, University of Wisconsin-Madison, 475 N. Charter St., Madison, WI 53706, USA.
 lazarian@astro.wisc.edu; esquivel@astro.wisc.edu

parallel to $\mathbf{r}$, respectively) 3D spectra of the velocity field, which are functions of the wavenumber amplitude k .

Similarly, the two point density statistics can be given by the structure function $\langle(\rho_1 - \rho_2)^2\rangle \equiv d(r)$.

3. MODIFIED VELOCITY CENTROIDS (MVCS)

In what follows we consider the emissivity of our media proportional to the first power of density² and no absorption present. In this case, the intensity, $I(\mathbf{X}) \equiv \int \rho_s dV$, is proportional to the column density, where ρ_s is the density of emitters in the Position-Position Velocity (PPV) space (see Lazarian & Pogosyan 2000). That is, $I(\mathbf{X}) = \int \rho_s dV = \int \alpha \rho_{tot} dz$, where, ρ_{tot} is the density of atoms, α is a proportionality constant, dz denotes integration along the line of sight, and $\mathbf{X}$ is a two dimensional vector in the plane of the sky.

The structure function of MVCs consists of two terms. It will be clear from the discussion below that it may be safe to use ordinary centroids when the second term is much smaller than the first term. In particular,

$$M(\mathbf{R}) = \left\langle [S(\mathbf{X}_1) - S(\mathbf{X}_2)]^2 - \langle V^2 \rangle [I(\mathbf{X}_1) - I(\mathbf{X}_2)]^2 \right\rangle \quad (3)$$

where $\mathbf{R} = \mathbf{X}_1 - \mathbf{X}_2$, $S(\mathbf{X})$ is a “unnormalized centroid” of the z -component of velocity V_z , and $\langle V_z^2 \rangle = v_0^2 + v_{turb}^2 + v_{thermal}^2$. To simplify our notations we shall omit the *turb* and z -subscripts (also for eq.(3)). We do not discuss the effect of $v_{thermal}$ separately because for the technique presented here it acts in the same way as regular motion, and one can formally introduce $v_0'^2 = v_0^2 + v_{thermal}^2$. $S(\mathbf{X}) \equiv \int_{\mathbf{X}} V \rho_s dV$, that is, at a given position $\mathbf{X}$, integrating along the line of sight³.

From an observational standpoint, the velocity dispersion $\langle V^2 \rangle$ can be obtained using the second moment of the spectral lines:

$$\langle V^2 \rangle \equiv \left\langle \int_{\mathbf{X}} V^2 \rho_s dV \right\rangle / \left\langle \int_{\mathbf{X}} \rho_s dV \right\rangle. \quad (4)$$

To express the 2D statistics of the modified centroids $M(\mathbf{R})$ through the underlying 3D velocity and density statistics we shall have to make several elementary transformations. First of all, the difference of integrals that enter the expression for the modified centroids should be presented as a double integral. Then using an elementary identity $(a-b)(c-d) = 1/2[(a-d)^2 + (b-c)^2 - (a-c)^2 - (b-d)^2]$ it is possible to find (see Lazarian 1995)

$$\langle (S(\mathbf{X}_1) - S(\mathbf{X}_2))^2 \rangle = \alpha^2 \iint dz_1 dz_2 [D(\mathbf{r}) - D(|z_1 - z_2|)], \quad (5)$$

where $D \equiv \langle (V_1 \rho_{tot,1} - V_2 \rho_{tot,2})^2 \rangle$. Using the Millionshikov hypothesis (see Monin & Yaglom 1975) to relate the fourth moments of the fields with the second moments, namely that $\langle h_1 h_2 h_3 h_4 \rangle \approx \langle h_1 h_2 \rangle \langle h_3 h_4 \rangle +$

$\langle h_1 h_3 \rangle \langle h_2 h_4 \rangle + \langle h_1 h_4 \rangle \langle h_2 h_3 \rangle$ we get sufficient simplifications of the expression:

$$D \approx \langle V^2 \rangle d(r) + \langle \rho_{tot}^2 \rangle \langle (V_1 - V_2)^2 \rangle - \frac{1}{2} \langle (V_1 - V_2)^2 \rangle d(r) + c(\mathbf{r}), \quad (6)$$

where $c(\mathbf{r})$ denote the cross terms arising from correlations between the fluctuations of the velocity and density

$$c(\mathbf{r}) = 2 \langle v_1 \rho_2 \rangle^2 - 4 v_0 \langle v_1 \rho_1 \rho_2 \rangle - 4 \rho_0 \langle \rho_1 v_1 v_2 \rangle. \quad (7)$$

The correlations between velocity and density fluctuations have been studied earlier (Esquivel et al. 2003; Lazarian et al. 2001) and found not to have a strong impact on the VCA. In particular, the first term in eq. (7) was directly obtained in Esquivel et al. (2003), and numerical integration of the corresponding fluctuations provides fluctuations in the centroid value that does not exceed 8% of the MVCs values at small $|R|$. In this paper we will neglect the cross terms and test this assumption by analyzing synthetic maps of compressible MHD simulations. A more detailed analysis, with more sets of data, and the cross terms, will be included in a forthcoming paper.

Eq. (6) depends on both density and velocity statistics. The dominant contribution arising from density fluctuations is the first term of eq. (6). Integration of this term yields the second term in eq. (3), as result the expression for the MVCs is

$$M(\mathbf{R}) \approx \alpha^2 \langle V^2 \rangle \langle \rho_{tot}^2 \rangle \iint dz_1 dz_2 [b(\mathbf{r}) - b(|z_1 - z_2|)], \quad (8)$$

where

$$b(\mathbf{r}) = \frac{\langle (V_1 - V_2)^2 \rangle}{\langle V^2 \rangle} \left(1 - \frac{1}{2} \frac{d(r)}{\langle \rho_{tot}^2 \rangle} \right). \quad (9)$$

If $(1/2)(d(r)/\langle \rho_{tot}^2 \rangle) \ll 1$ then eq.(9) recovers the statistics of the velocity field. This is certainly the case if the turbulence is subsonic, where the amplitude of the density fluctuations is small, or when the density spectrum is steep (i.e. most of the power residing at large scales) given that we are interested in the fluctuations at small R .⁴ However, if the amplitude fluctuations of density at small scales is large (i.e. shallow spectrum or highly supersonic turbulence), the contribution could be important. If we omit the density term in eq.(9) and take a 2D Fourier transform to resulting distribution of MVCs we have:

$$P_2(\mathbf{K}) = \frac{1}{4\pi^2} \int d^2\mathbf{R} e^{-i\mathbf{K}\cdot\mathbf{R}} M(\mathbf{R}), \quad (10)$$

where we have used again the notation of using capital letters to distinguish 2D vectors from the 3D counterparts. Substituting eqs. (8), (1), and (2) into eq. (10) one gets

$$P_2(K) \sim C_1 F_{NN}(K, 0) + C_2 \delta(K) \quad (11)$$

where C_1 and C_2 are constants ($C_1 = 4\pi\alpha^2 \langle \rho_{tot}^2 \rangle L$, where L is the extent of the emitting region) and the second term provides only a δ function contribution at $K = 0$. Thus eq. (11) provides a relation⁵ between the spectrum of the modified centroids and the

² This is the case for the H I 21cm line. For recombination radiation (such as H α for instance) the emissivity is proportional to the square of the density, and modifications of the present technique are necessary.

³ In terms of practical data handling, to minimize the effect of absorption at the centers of the lines, subtracting of mean values of velocity v_0 in the expression for $S(\mathbf{X})$ and $\langle V^2 \rangle$ may be advantageous. In terms of $S(\mathbf{X})$ the use of $\delta S(\mathbf{X}) \equiv S(\mathbf{X}) - \langle S(\mathbf{X}) \rangle$ may be preferable.

⁴ By *steep* spectrum we mean $P_{3D} \propto k^n$ with $n < -3$. If $n > -3$ the spectrum is *shallow* (see Lazarian & Pogosyan 2000).

⁵ Our result is similar to that in Kitamura (1993) who used normalized centroids given by eq. (12) but implicitly assumed that $\rho = \text{const}$ (see also Vestuto et al. 2003).

transversal spectrum of the 3D velocity field. We should note that the relation in eq. (11) does not depend on a particular form (power-law is a common assumption) of the underlying spectra. If the turbulent velocity field is mostly solenoidal (see Matthaeus et al. 1996; Porter, Woodward & Pouquet 1998), the isotropic spectrum $E(k)$ that is used for describing hydrodynamic turbulence is uniquely defined through $F_{NN}(k)$, namely, $E(k) \approx 4\pi k^2 F_{NN}(k)$.

4. NUMERICAL TESTING

In what follows we will include the normalized centroids in their usual form (see Miesch & Bally 1994)

$$C(\mathbf{X}) \equiv \int_{\mathbf{X}} V \rho_s dV / \int_{\mathbf{X}} \rho_s dV. \quad (12)$$

Intuitively it is clear that the power spectrum of the centroids given by eq. (12) should provide a better fit to the power spectrum of the velocity (the spectral index, not the amplitude) than that of $S(\mathbf{X})$. Indeed, the contributions of density fluctuations are mitigated by the division over the column density in eq. (12).

The testing was done using the files obtained through simulations of compressible MHD turbulence. The description of the code and the numerical simulations can be found in Cho & Lazarian (2002) and in Esquivel et al. (2003). We use the same data cube that we used in Esquivel et al. (2003). Namely, the dimension of the cube is 216^3 , and the Mach number 2.5. The result of the MHD simulations are velocity, density and magnetic field data. We use density and velocity to produce spectral line data cubes, the procedure is described in Esquivel et al. (2003).

We calculated $S(\mathbf{X})$, $C(\mathbf{X})$, and $I(\mathbf{X})$ from synthetic spectral line data cubes. The spectra $P_2(\mathbf{K})$ were obtained by applying a fast Fourier transform to the 2D distribution of the centroids and column density values. The power spectrum of the MVCs was calculated subtracting $\langle V^2 \rangle$ times the column density power spectrum, from the power spectrum of $S(\mathbf{X})$ (see eq. 3). For the original data cubes (with a steep density spectrum), the differences among all three types of centroids were marginal for the dynamical range studied, although unnormalized centroids systematically show more deviations from the velocity power spectrum. To enhance the effect of density, we made the density shallow by changing the Fourier components of the density amplitudes, but, in order to preserve the density-velocity correlations, we kept the phase information (see Esquivel et al. 2003). The results are shown in Fig 1. It is clear that the three measures provide a good fit at large scales, but at small scales normalized and unnormalized centroids tend to be density dominated. An additional advantage of using MVC over the conventional normalized centroids is that MVC provides the correct amplitude of the spectrum, while normalized centroids at best give the correct index (logarithmic slope).

It is also clear that for a shallow density spectrum, the improvement from unnormalized centroids to normalized centroids is marginal.

Our numerical study shows that MVCs are advantageous when the underlying spectrum of density is dominated by fluctuations at small scales. And, the main




FIG. 1.— The comparison of the 2D spectrum of the three types of centroids with the underlying 3D spectrum of velocity fluctuations. The solid line corresponds to the power spectrum of velocity for compressible MHD simulations with Mach number ~ 2.5 . The dotted line corresponds to the spectrum of density modified by changing the amplitudes the Fourier components of the original spectrum. The modified density spectrum $E(k)/k^2 \sim k^{-2.5}$ for the inertial range. The vertical position of the normalized centroids (triangles) is shifted vertically for visual purposes.

advantage of MVCs is that they allow an explicit statistical description of the procedures involved, which makes the analysis more reliable.

5. DISCUSSION

Our study revealed that the MVCs can be successfully used to obtain the velocity statistics from synthetic observational data. Another technique, namely, the VCA is complementary to the MVCs. VCA is more robust to density-velocity correlations. For instance, it was shown in Lazarian & Pogosyan (2000) that VCA can provide correct statistics of velocity fluctuations even in the case when the velocity and density are correlated at their maximum given by the Cauchy-Swartz inequality (see Mathews & Walker 1970). On the contrary, if we assume such a maximum correlation for the MVCs we will not get the correct scaling. Comparing results obtained with the MVCs and the VCA one can get a better handle on what are the velocity statistics.

It is advantageous to combine VCA and MVCs techniques. We can apply VCA to the largest scales, where turbulence is supersonic, and study MVCs at all scales, including those where the turbulence becomes subsonic. The correspondence of the spectral slope obtained with the two different techniques would substantially increase the reliability of the result. There is a more subtle, but important point. MVCs provide the spectrum of solenoidal motions given by F_{NN} , while VCA is sensitive to both potential and solenoidal motions. This may be used to get both potential and solenoidal motions in order to estimate the role of compressibility. In addition,

because the thermal velocity acts in the VCA analysis the same way as the thickness of the channel maps, measurements of the spectrum with the MVCs can provide a means of estimating the thermal velocity. It is essential to combine the advantages of the techniques.

Our analysis of numerical data in §4 shows that the worries about the velocity centroids (see Stenholm 1990) although justified, may be somewhat exaggerated.

If we *assume* that the criterion for the use of the centroids is satisfied (that the second term in eq. (3) is small compared to the first one) for earlier data sets, the analysis of observational data in (Miesch & Bally 1994) provides a range of power-law indexes. Their results obtained with structure functions if translated into spectra are consistent with $E(k) = k^\beta$, where $\beta = -1.86$ with a standard deviation of 0.3. The Kolmogorov index $-5/3$ falls into the range of the measured values. L1228 exhibits exactly the Kolmogorov index -1.66 as the mean value, while other low mass star forming regions L1551 and HH83 exhibit indexes close to those of shocks, i.e. ~ -2 . The giant molecular cloud regions show shallow indexes in the range of $-1.9 < \beta < -1.3$ (see Miesch, Scalo & Bally 1999). It worth noting that Miesch & Bally (1994) obtained somewhat more shallow indexes that are closer to the Kolmogorov value using autocorrelation functions. Those may be closer to the truth as in the presence of absorption in the center of lines, minimizing the regular velocity used for individual centroids might make the results more reliable (see also first footnote in §3). Repeating of the study using MVCs and taking care of the absorption effects may be advantageous.

The emergence of the Kolmogorov index for the compressible magnetized gas may not be so surprising. Cho & Lazarian (2002) for a range of Mach numbers reported the existence of distinct Alfvénic, slow, and fast

wave cascades. The Alfvén and slow waves result in Kolmogorov-type scaling. The application of the VCA to Small Magellanic Cloud (Stanimirović & Lazarian 2001) resulted in the velocity spectrum consistent with those predictions. However, other researchers (e.g. Vestuto et al. 2003) reported indexes that varied with the Mach number of the simulations. It is clear that more theoretical and observational work is necessary.

6. SUMMARY

1. We derived a criterion when the velocity centroids may reflect the actual underlying velocity statistics. We introduced MVCs that may recover velocity statistics from spectral line data in cases when the traditional centroid analysis fails.

2. Our numerical tests show that both the MVCs and the normalized velocity centroids (see eq. (12)) successfully recover the velocity information from data cubes obtained via numerical simulations of the compressible MHD turbulence if the density spectrum is steep.

3. MVCs and the VCA are complementary independent techniques. Obtaining the same power spectrum with both of them enhances confidence in the result. MVCs are most useful when turbulence is subsonic or/and does not follow the power law. Additional information on the solenoidal versus potential motions, temperatures of gas etc. may be obtained combining these two techniques.

We thank the referee Anthony Minter for valuable suggestions that improved this work. The research by A.L. is supported by the NSF Grant AST-0125544. A.E. acknowledges financial support from CONACyT (Mexico). Fruitful communications with Volker Ossenkopf are acknowledged.

REFERENCES

- Cho, J., & Lazarian, A. 2002, Phys. Rev. Lett., 88, 5001
 Cho, J., & Lazarian, A. 2003a, in Acoustic emission and scattering by turbulent flows, ed. M. Rast (Springer LNP), in press, astro-ph/0301462
 Cho, J., & Lazarian, A. 2003b, MNRAS, submitted, astro-ph/0301062
 Deshpande, A. A., Dwarakanath, K. S., Goss, W. M. 2000, ApJ, 543, 227
 Esquivel A., Lazarian A., Pogossyan D., Cho J. 2003, MNRAS, in press, astro-ph/0210159
 Kitamura, Y., Sunada, K., Hayashi, M., Hasegawa, T. 1993, ApJ, 413, 221
 Larson R. B., 1992, MNRAS, 641
 Lazarian, A. 1995, A&A, 293, 507
 Lazarian, A. 1999, in Plasma Turbulence and Energetic Particles in Astrophysics, eds. Michał Ostrowski, Reinhard Schlickeiser, Kraków 1999, 28, astro-ph/0001001
 Lazarian, A., Pogossyan, D. 2000, ApJ, 537, 72
 Lazarian, A., Pogossyan, D., Vázquez-Semadeni, E., Pichardo, B. 2001, ApJ, 555, 130
 Matthaeus, W. M., Ghosh, S., Oughton, S., Roberts, D. A. 1996, J. Geophys. Res., 101, 7619
 Mathews, J., & Walker, R.L. 1970, Mathematical Methods of Physics, NY: Benjamin
 Miesch, M. S., Bally, J. 1994, ApJ, 429, 645
 Miesch, M. S., Scalo, J., & Bally, J. 1999, ApJ, 429, 645
 Monin, A. S., Yaglom, A. M. 1975, Statistical fluid dynamics: mechanics of turbulence, Vol. 2., MIT press, Cambridge, MA
 Munch, G., 1958, Rev. Mod. Phys., 30, 1035
 Porter, D., Woodward, P., Pouquet, A. 1998, Phys. of Fluids, 10, 237
 Spoelstra, T. A. T. 1972, A&A, 21, 61
 Stanimirović, S., & Lazarian, A., 2001, ApJ, 551, 53
 Stenholm, L.G. 1990, A&A, 232, 495
 Vázquez-Semadeni, E., Ostriker, E.C., Passot, T., Gammie, C.F., & Stone, J.M. 2000, in Protostars and Planets IV, eds. V. Mannings et al. (Tucson: University of Arizona Press), p.3
 Vestuto, J.G., Ostriker, E.C., & Stone, J.M. 2003, ApJ, in press, astro-ph/0303103